

## Energy analysis and CO<sub>2</sub> eq emissions of chicken meat production

### Análisis energético y emisiones de CO<sub>2</sub> eq en la producción de carne de pollo

Eduardo Baltierra-Trejo<sup>1</sup>, Alejandra Arroyo-Pitacua<sup>2</sup>, Liliana Márquez-Benavides<sup>2\*</sup>

<sup>1</sup>Centro de Investigación en Sustentabilidad Energética y Ambiental del Noreste, Universidad Autónoma del Noreste, CP. 87560. Matamoros, Tamaulipas, México.

<sup>2</sup>Laboratorio de Residuos Sólidos y Medio Ambiente del Instituto de Investigaciones Agropecuarias y Forestales. Universidad Michoacana de San Nicolás de Hidalgo. CP. 58302. Morelia, Michoacan, México.

\*Autor de correspondencia: lmarquez@umich.mx

Scientific note received: May 11, 2016 accepted: November 23, 2016

**ABSTRACT.** In México, greenhouse gas emissions from the agricultural sector include mainly enteric and manure sources. However, emissions due to the use of fossil energy are usually not considered. The aim of this work was to identify the energy demand (MJ) to produce a kilogram of chicken meat, and to determine the associated CO<sub>2</sub> equivalent emissions. To that end, a farm in west-central Mexico was studied to profile the energy demand to supply water, feed, lighting, ventilation, air extraction and heating, using 1000 birds as a calculation basis. It was found that the emissions derived from fossil energy use were 0.47 kgCO<sub>2</sub> eq per kilogram of live weight per production cycle.

**Key words:** Electricity, energy, energy efficiency, poultry

**RESUMEN.** En México, las emisiones de gases de efecto invernadero del sector agropecuario son principalmente entéricas y de manejo de estiércol, pero no se consideran las emisiones de la demanda de energética fósil. Con el objetivo de conocer la demanda energética (MJ) para producir un kilogramo de carne de pollo y determinar la emisión de CO<sub>2</sub> equivalente asociada, se analizó el consumo energético necesario para el suministro de agua, alimento, iluminación, ventilación, extracción de aire y calefacción en una granja, tomando como base de cálculo 1000 aves. Se determinó que la emisión derivada del consumo de energía fósil fue de 0.47 kg CO<sub>2</sub> eq por kilogramo de carne de pollo en pie en un ciclo productivo.

**Palabras clave:** Electricidad, energía, eficiencia energética, avícola

## INTRODUCTION

The agricultural sector accounts for an estimated 20 % of global greenhouse gas (GHG) emissions (IPCC 2014). Mexico ranks 12th in relation to CO<sub>2</sub> equivalent (CO<sub>2</sub> eq) emissions with 374 million tons, of which 12.3 % comes from agricultural activities (SEMARNAT and INECC 2012). Nationally, the poultry industry accounts for 63 % of livestock production, but the national inventory of greenhouse gas emissions only reports CO<sub>2</sub> emissions for manure management in the general agriculture category. Therefore, the carbon footprint for fossil energy use by the poultry activi-

ty is unknown (SEMARNAT and INECC 2013).

The Farm Energy Act (SAGARPA 2003) provides incentives in the energy rates for farm activities, in order to achieve greater productive efficiency. However, poultry farms require energy to: provide adequate thermal comfort, ventilation, and lighting to birds; mobilize feed, equipment, inputs and waste; dispose of mortalities and obtain water (Costantino *et al.* 2016). To determine the environmental impact of the energy consumption resulting from poultry farming, the matter and energy inputs and outputs from the chicken meat production process should be established and the CO<sub>2</sub> emissions generated should be estimated. Therefore, the ob-

jective was to define the energy demand (MJ) to produce one kilogram of chicken meat and to determine the associated CO<sub>2</sub> eq emissions.

## MATERIALS AND METHODS

### Location of the study area

The study was conducted in the summer of 2013 in a farm located in the municipality of Taretan, in the western region of the State of Michoacán, Mexico, located at 19° 20' 00" NL and 101° 55' 00" WL. The climate is temperate with summer rains, average annual rainfall of 1 560 mm and temperatures between 4.4 and 29.6 °C. The farm has 10 poultry houses of 13 x 150 m, with capacity for 22 000 birds per 49-day cycle. The production system is technified, with a controlled environment composed of heaters, fans, extractors, sprinklers for moisture control and a curtain system on the house walls.

### Determination of energy consumption

A survey conducted to learn the farm's overall process was applied to the poultry house keepers and the farm manager. The objective of the survey was to determine: the inputs and outputs of the process; the cycle's duration; consumption of LP gas (kg), electricity (kWh), feed (kg) and water (L); number of birds and live and carcass weight (kg) at the end of the production cycle. An inventory of machinery and equipment was carried out to determine the energy demand and hours of use. Fuel consumption in machinery and transportation vehicles was obtained from the farm's usage log.

The considerations for the calculations were that the basis of the report was for 1 000 birds during a 49-d production cycle; 13-hour d<sup>-1</sup> operation of the ventilation and air extraction systems; the use of heating and lighting was for 28 d during the 49-d production cycle; 1.7 kg carcass weight without head, neck, legs and viscera; and live weight of 2.54 kg at the end of the 49-d production cycle.

### Analysis of energy performance

It was determined with the energy productivity and specific energy equations of Singh (1997) and Salazar *et al.* (2012), which are as follows:

$$EP = \frac{\text{Production (kg, broiler live weight)}}{\text{Total energy consumed in a productive cycle (MJ)}}$$

Where EP = Energy productivity.

$$SE = \frac{\text{Total energy consumed in a productive cycle (MJ)}}{\text{Production (kg, broiler live weight)}}$$


Where SE = Specific energy.

### Determination of GHG emissions by energy consumption

GHG emissions from electric energy consumption to provide drinking water, feed, lighting, ventilation and air extraction were estimated based on the CO<sub>2</sub> eq emission factor for Mexico of 454 g CO<sub>2</sub> eq per kWh proposed by the International Energy Agency (IEA 2015). The National Institute of Ecology's carbon calculator (INECC and SEMARNAT 2013) was used to determine the GHG emissions of the LP gas-based heating system, while the determination of the electric energy consumption (kWh) used to extract and distribute drinking water (m<sup>3</sup>) was performed with the Michoacán efficiency index, which is 0.75 kWh m<sup>-3</sup> (CONUEE 2011).

## RESULTS AND DISCUSSION

The main system inputs and outputs for a broiler production cycle are described in Figure 1. The consumption of feed, water and electricity has variations throughout the production cycle, according to the age of the bird. The energy demand for the Taretan farm is summarized in Table 1; for 1 000 broilers the demand is 22 834 MJ. Therefore, the estimated energy productivity for broiler live weight is 0.12 kg MJ<sup>-1</sup>, while the specific energy demanded is 9.2 MJ kg<sup>-1</sup>. About 98 % of the energy consumption is due to the heating



**Figure 1.** Main inputs for the production of 1 000 broilers on a farm in the Municipality of Taretan in a 49-d production cycle.

**Table 1.** Supply of inputs and energy demanded for the production of 1 000 broilers at a farm in the Municipality of Taretan in a 49 d production cycle.

Time	Feed consumption		Water consumption		Lighting h week <sup>-1</sup>	Ventilation		Air extraction		Heating		
	kg	MJ	m <sup>3</sup>	MJ		MJ	kWh	MJ	kWh	MJ	kg LP gas	MJ
1	140	0.77	0.2	0.7	1.9	7.1	3.1	11.3	10.3	55	115.6	5 708.5
2	330	0.77	0.6	1.7	1.4	5.3	3.1	11.3	10.3	55	115.6	5 708.5
3	580	0.77	1.1	3.1	0.09	3.5	3.1	11.3	10.3	55	115.6	5 708.5
4	670	0.77	1.3	3.6	0.49	1.7	3.1	11.3	10.3	55	115.6	5 708.5
5	980	0.77	1.9	5.2	0	0	3.1	11.3	10.3	55	0	0
6	1 250	0.77	2.5	6.7	0	0	3.1	11.3	10.3	55	0	0
7	2 240	0.77	4.4	12.0	0	0	3.1	11.3	10.3	55	0	0
Total per category	6 190	5.4	12.4	33.4	4.9	17.8	22	79	72	385	462	22 834
Cycle total = 23 354 MJ												
GHG emissions (kgCO <sub>2</sub> eq)		2.4	4		2.2		10		33		1 157	
Cycle total = 1 208.6 kgCO <sub>2</sub> eq												

system, even though LP gas is required in only four weeks of the production cycle. This coincides with the figure reported by Sonesson *et al.* (2009), who indicate that fuel consumption for heating accounts for between 80 and 90 % of the total energy used, which suggests an area of opportunity in which strategies that reduce energy consumption should be focused on. On the other hand, an FAO study (2013) found that the energy con-

sumption was 4.5 MJ kg<sup>-1</sup> carcass weight, which is lower than that found in the present study, which may be due to the fact that this study took into account the energy consumed by the ventilation and heating systems. In addition, the energy productivity for broiler live weight of 0.12 kg MJ<sup>-1</sup> is higher than that obtained in a simulation carried out by Sefat *et al.* (2014), where they obtained an energy productivity of 0.01 kg MJ<sup>-1</sup>, in which the

**Table 2.** Specific energy for the production of chicken meat, pork and beef.

Category	Poultry carcass MJ kg <sup>-1</sup>	Pork MJ kg <sup>-1</sup>	Beef MJ kg <sup>-1</sup>
Electricity			
Water supply	0.01	0.02	-
Lighting	0.01	0.05	-
Feed supply	0.002	0.02	-
Ventilation	0.03	0.20	-
Air extraction	0.15	-	-
High-pressure wash	-	0.03	-
Auxiliary heating	-	0.19	-
Subtotal	0.20	0.51	2.34
LP gas heating	8.96	0.81	-
Diesel	-	-	7.92
Manure management	-	0.02	-
Hot water	-	0.12	-
Total	9.2	1.45	10.26
Source	This study	Lammersa <i>et al.</i> (2010)	Sonesson <i>et al.</i> (2009)

**Table 3.** CO<sub>2</sub> eq emissions from different chicken meat production processes.

Source	kg CO <sub>2</sub> eq kg <sup>-1</sup> of chicken meat	kg CO <sub>2</sub> eq kg <sup>-1</sup> of pork carcass	kg CO <sub>2</sub> eq kg <sup>-1</sup> of beef carcass
This study	0.47 $\diamond$	-	-
This study	0.7 $\Delta$	-	-
Roy <i>et al.</i> (2011)	4.57 $\diamond$	5.57	34.3
Cederberg <i>et al.</i> (2009)	1.93 $\Delta$	3.4	19.8
Thynelius (2008)	1.5 $\Delta$	-	-
Williams <i>et al.</i> (2006)	7.3 $\Delta$	-	-
Wiedemann <i>et al.</i> (2017)	2.2 $\ddagger$	-	-

$\diamond$  Live weight,  $\Delta$  Broiler carcass weight,  $\ddagger$  C Frozen Chicken Meat .


energy contributions for feed production and manure management were considered. In comparing the specific energy required for the production of chicken meat (Lammersa *et al.* 2010) and beef (Sonesson *et al.* 2009), they were found to have similar values (Table 2), but they come from different inputs, such as consumption of LP gas for heating chickens and the consumption of diesel for the machinery required for beef production.

The energy demand translates into 1 206.4 kg CO<sub>2</sub> eq (Table 1). That is, 0.47 kg CO<sub>2</sub> eq are generated per kilogram of broiler live weight or 0.70 kg CO<sub>2</sub> eq to produce a kilogram of carcass meat. The farm's CO<sub>2</sub> eq emissions (Table 3) were lower than those of other poultry systems (Thynelius 2008, Williams *et al.* 2006, Wiedemann *et al.* 2017) and those reported for the production of pork and beef (Roy *et al.* 2011, Cederberg *et al.* 2009). However, it is necessary to take into account the following considerations when comparing CO<sub>2</sub> eq emissions between livestock production systems: 1)

differences in production systems evaluated, climatic conditions and management practices of each farm; 2) the limits of research in each report vary, and 3) some results were calculated on the basis of estimates made in other countries, such as that of Roy *et al.* (2011) made for meat production in Japan and that of Cederberg *et al.* (2009) in Sweden.

The present study focuses on the energy consumption inside the poultry house, but does not consider the production of poultry feed, a process reported as the largest energy consumer (Nguyen *et al.* 2012, Wiedemann *et al.* 2017). The energy used during the production of poultry feed was calculated by Nguyen *et al.* (2012); if this value is added to the figure already obtained, the estimated GHG emissions per kilogram of broiler live weight increases to 61.9 kgCO<sub>2</sub> eq. According to Pelletier (2008), feed provision accounts for approximately 80 % of the energy consumed and 82 % of greenhouse gas emissions.

There are no national estimates for GHG



**Figure 2.** Estimation of the contribution of poultry farming to GHG emissions in Mexico: Agriculture Subcategory.

emissions from poultry farming in Mexico. However, if the farm studied is considered to be typical in this country, the results suggest that the GHG contribution by poultry farms would increase the emissions reported for the agriculture subcategory by 13 % (SEMARNAT and INECC 2013) (Figure 2).

Energy demand for the production of one kilogram of chicken meat and the associated CO<sub>2</sub> eq emissions were determined; this information can serve as a starting point to know the average actual energy consumption of the poultry industry in

Mexico and to find areas of opportunity to reduce GHG emissions. The next step should be to determine the energy demand of the entire supply chain to know the total energy consumption for the production of chicken meat and its derivatives.

## ACKNOWLEDGEMENTS

The authors are grateful for Project 2014-2015 CIC-UMSNH funding and to CONACYT for postgraduate scholarship 23918.

## LITERATURA CITADA

- Cederberg C, Flysjö A, Sonesson U, Sund V, Davis J (2009) Greenhouse gas emissions from Swedish consumption of meat, milk and eggs 1990 and 2005. SIK Report 793. The Swedish Institute for Food and Biotechnology. Estocolmo, Suecia. 96p.
- Costantino A, Fabrizio E, Biglia A, Cornale P, Battaglini L (2016) Energy use for climate control of animal houses: The state of the art in Europe. Energy Procedia 101: 184-191.
- CONUEE (2011) Estudio Integral de Sistemas de Bombeo de Agua Potable Municipal. Comisión Nacional para el Uso Eficiente de Energía - Secretaría de Energía, Cooperación Alemana al Desarrollo. México. 98p.

- IEA (2015) CO<sub>2</sub> emissions from fuel combustion Highlights. The International Energy Agency. Paris, France. <https://www.iea.org/publications/freepublications/publication/CO2EmissionsFromFuelCombustion-Highlights2015.pdf>. 139p. Date consulted: April 19, 2016.
- IPCC (2014) Cambio climático 2014: Informe de síntesis. Contribución de los Grupos de trabajo I, II y III al quinto informe de evaluación del grupo intergubernamental de expertos sobre el cambio climático. Intergovernmental Panel on Climate Change. Ginebra, Suiza. 157p.
- Lammersa PJ, Honeymana MS, Harmonb JD, Helmersb MJ (2010) Energy and carbon inventory of Iowa swine production facilities. *Agricultural Systems* 103: 551-561.
- MacLeod M, Gerber P, Mottet A, Tempio G, Falcucci A, Opio C, et al. (2013) Greenhouse gas emissions from pig and chicken supply chains, A global life cycle assessment. Food and Agriculture Organization of The United Nations Animal Production and Health Division. Roma, Italia. 170p.
- Nguyen TTH, Bouvarel I, Ponchant P, Hayo MG, Werf VD (2012) Using environmental constraints to formulate low-impact poultry feeds. *Journal of Cleaner Production* 28: 215-224.
- Pelletier N (2008) Environmental performance in the US broiler poultry sector: life cycle energy use and greenhouse gas, ozone depleting, acidifying and eutrophying emissions. *Agricultural Systems* 98: 67-73.
- Roy P, Orikasa T, Thammawong M, Nakamura N, Xu Q, Shiina T (2011) Life cycle of meats: an opportunity to abate the greenhouse gas emission from meat industry in Japan. *Journal of Environmental Management* 93: 218-224.
- SAGARPA (2003) Ley de Energía Para el Campo. Diario Oficial de la Federación el 4 de diciembre de 2003. Ciudad de México, México. 5p. [http://www.diputados.gob.mx/LeyesBiblio/regley/Reg\\_LECampo.pdf](http://www.diputados.gob.mx/LeyesBiblio/regley/Reg_LECampo.pdf). Date consulted: April 19, 2016.
- Salazar M, Cruz R, Rojano P (2012) Eficiencia en el uso de la energía en invernaderos mexicanos. *Revista Mexicana de Ciencias Agrícolas* 4: 736-742.
- Sefat M, Borgaee A, Beheshti B, Bakhoda H (2014) Modeling energy efficiency in broiler chicken production units using artificial neural network (ANN). *International Journal of Natural and Engineering Sciences* 8: 7-14.
- SEMARNAT, INECC (2013) Inventario nacional de emisiones de gases de efecto invernadero 1990-2010. Secretaría de Medio Ambiente y Recursos Naturales, Instituto Nacional de Ecología y Cambio Climático. México. 385p.
- SEMARNAT y INECC (2012) México quinta comunicación nacional ante la Convención Marco de las Naciones Unidas Sobre el Cambio Climático. Instituto Nacional de Ecología. Secretaría de Medio Ambiente y Recursos Naturales. Instituto Nacional de Ecología y Cambio Climático. México. 396p.
- Singh MK, Pal SK, Thakur R, Verma UN (1997) Energy input-output relationship of cropping systems. *Indian Journal of Agricultural Science* 67: 262-266
- Sonesson U, Cederberg C, Berglund M (2009) Greenhouse gas emissions in chicken production. Decision support for climate certification. *Klimatmärkning För Mat*, Report 2009:6. Estocolmo, Suecia. 19p. <http://www.klimatmarkningen.se/wp-content/uploads/2009/12/2009-6-chicken.pdf>. Date consulted: April 19, 2016.
- Thynelius G (2008) Klimatpåverkan och förbättringsåtgärder för Lantmännens livsmedel - fallstudie Kronfågels slaktkyckling. Examensarbete 2008. Institutionen för teknik och samhälle, Miljö- och Energisystem, Lunds tekniska högskola, Lund. Lund, Suecia. 50p.

- Wiedemann SG, McGaham EJ, Murphy CM (2017) Resource use and environmental impacts from Australian chicken meat production, *Journal of Cleaner Production* 140: 675-684.
- Williams AG, Audsley E, Sandars DL (2006) Determining the environmental burdens and resource use in the production of agricultural and horticultural commodities, Main Report, Defra Research project ISO205. Cranfield University och Defra. Bedford, Inglaterra. 46p.

